

Bend, Break and Count II: elliptics, cuspidals, linear genera

Z. Ran

Department of Mathematics
University of California, Riverside
Riverside, CA 92521 USA
ziv@ucrmath.ucr.edu

In [R2] we showed how elementary considerations involving geometry on ruled surfaces may be used to obtain recursive enumerative formulae for rational plane curves. Here we show how similar considerations may be used to obtain further enumerative formulae, as follow.

Theorem. (i) *The number N_d^1 of elliptic plane curves of degree d through $3d$ general points satisfies*

$$(1) \quad N_d^1 = \frac{1}{12} \binom{d}{3} N_d^0 + \sum_{d_1+d_2=d} \frac{3d_1-2}{9} d_1 d_2 N_{d_1}^0 N_{d_2}^1 \binom{3d-1}{3d_1-1};$$

(ii) *The numbers K_d^0, K_d^1 of rational (resp. elliptic) plane curves with 1 cusp through $3d-2$ (resp. $3d-1$) general points satisfy*

$$(2) \quad K_d^0 = 3N_d^0 - \sum_{d_1+d_2=d} N_{d_1}^0 N_{d_2}^0 d_1 d_2 \left[(3d_2-2) \binom{3d-2}{3d_1-2} - \frac{3}{2} \binom{3d-4}{3d_1-2} \right],$$

$$(3) \quad K_d^1 = 3N_d^1 + \frac{(d-1)(d-2)(d-4)}{8} N_d^0 + \sum (3d_1-2) d_1 d_2 \binom{3d-1}{3d_1-1} N_{d_1}^0 N_{d_2}^1;$$

(iii) *the respective linear genera (= geometric genera of generic curve-sections) g_d^0, g_d^1 of the Severi varieties V_d^0, V_d^1 , satisfy*

$$(4) \quad 2g_d^0 - 2 = K_d^0 - \sum N_{d_1}^0 N_{d_2}^0 d_1 d_2 \binom{3d-4}{3d_1-2},$$

$$(5) \quad 2g_d^1 - 2 = K_d^1 - \frac{9}{2} N_d^1 + \frac{(d-1)(d-2)(3d-4)}{24} N_d^0 + \sum \frac{3d_1-2}{2} d_1 d_2 N_{d_1}^0 N_{d_2}^1 \binom{3d-1}{3d_1-1}.$$

Here (1) is due to Getzler and Pandharipande [P] while the other formulae seem new.

Acknowledgement. This work was begun in June '97 under the auspices of the 'Trimestre di geometria numerativa' at the 'Università di Roma Tor Vergata'. I am grateful to the Trimestre's organizers, Ciro Ciliberto and Corrado De Concini, as well as other participants, for the stimulating and congenial atmosphere, and to Rahul Pandharipande for telling me about his recent work.

Now let us denote, for any function u of d_1 ,

$$T(u) = \sum_{d_1+d_2=d} u(d_1) d_1 d_2 \binom{3d-1}{3d_1-1} N_{d_1}^0 N_{d_2}^1.$$

Then as each R_i of degree d_1 contains precisely $3d_1 - 1$ of the A 's, we have

$$s_{A_i}^2 = -\omega - \frac{1}{3d-1} T(3d_1 - 1).$$

Now consider the ramification divisor of the natural map

$$f : X \rightarrow \mathbb{P}^2.$$

This contains a vertical part $\sum_1^\kappa F_i$ coming from the cuspidal curves. Along a smooth fibre $F = \pi^{-1}(b)$, the ramification points of f are given by the residual intersection of $f(F)$ with the unique (mod $f(F)$) plane curve C of degree d through the nodes of $f(F)$, which intersection is given by $A_1, \dots, A_{3d-1}$ plus one further point $\sigma(b)$. For a singular fibre F the situation is similar except C is only required to pass through the nodes of $f(F)$ not coming from the node of F ; moreover, it is easy to see in case $F = E_i + R_i$ that the 'last' point $\sigma(b)$ lies on $f(E_i)$, the elliptic part. A moment's thought shows that the curves $f(F)$ having a node at some A_i do not require special consideration; what happens there is that $\bar{B}$ itself will have a node at the point $\{f(F)\}$, with branches corresponding to the branches of $f(F)$ through A_i and $s_{A_i}(b)$ will be the point of F above A_i lying on the branch corresponding to the branch of $\bar{B}$ containing b . In any event, the points $\sigma(b)$ glue together to a section σ with

$$\begin{aligned} \sigma &\sim f^*(-K_{\mathbb{P}^2}) + K_X - \sum_1^{3d-1} s_{A_i} - \sum_1^\kappa F_i \\ &\sim 3L + (\omega + 2g - 2 - \kappa)F - \sum s_{A_i} + R \end{aligned}$$

where $L = f^*\mathcal{O}_{\mathbb{P}^2}(1)$, so that $L^2 = N_d^1$. As $\sigma \cap R = \emptyset$, we have

$$\begin{aligned} \sigma^2 &= -\omega \\ &= (3L + (\omega + 2g - 2 - \kappa)F - \sum s_{A_i} + R)^2. \end{aligned}$$

Simplifying, we get

$$(8) \quad 2g - 2 - \kappa = \frac{1}{2}((3d-9)\omega - gN_d^1 + T(3d_1-2))$$

On the other hand, we have (as all the s_{A_i} are interchangeable)

$$\begin{aligned} (\sigma + \sum s_{A_i})^2 &= -3d\omega - T(3d_1-1) + 2(3d-1)\sigma \cdot s_{A_1} \\ &\quad + (3L + (\omega + 2g - 2 - \kappa)F - R)^2 \end{aligned}$$

Simplifying, we get

$$(9) \quad \sigma.s_{A_1} = \frac{1}{2(3d-1)}(9N_d^1 + 9d\omega + 6d(2g-2-\kappa) + T(9d_1-2)).$$

Now we compute $\sigma.s_{A_1}$ another way. The key to this is the observation that the intersections $\sigma \cap s_{A_1}$ correspond precisely to those fibres F on which the line bundle $\mathcal{O}_X(3L+R+\pi^*M)$, for any divisor M on B , has the ‘same’ (i.e. isomorphic) restriction as $\mathcal{O}_X(2s_{A_1} + s_{A_2} + \dots + s_{A_{3d-1}})$; a moment’s thought shows that this is valid even for the singular fibres and that, by general choice, the intersection is transverse. To make use of this observation, consider the natural restriction map

$$\rho : V := \pi_*(\mathcal{O}_X(3L+R)) \rightarrow W := \pi_*(\mathcal{O}_{2s_{A_1}+s_{A_2}+\dots+s_{A_{3d-1}}}(3L+R)).$$

Then V and W both have rank $3d$ and $\pi(\sigma \cap s_{A_1})$ coincides with the degeneracy locus of ρ , hence

$$(10) \quad \sigma.s_{A_1} = c_1(W) - c_1(V).$$

Now it is easy to see that

$$(11) \quad c_1(W) = \omega + \frac{3d+1}{3d-1}T(3d_1-1).$$

As for $c_1(V)$, it may be computed from Grothendieck-Riemann-Roch [F], which yields

$$(12) \quad \begin{aligned} c_1(V) + 3d(1-g) &= ((1 + (3L+R) + \frac{1}{2}(3L+R)^2) \cup (1 - \frac{K_X}{2} + \chi(\mathcal{O}_X)[pt]))([X]) \\ &= \frac{(3L+R)^2}{2} - \frac{K_X}{2}(3L+R) + \omega \end{aligned}$$

Now (12) yields a formula for $c_1(V)$, from which the g term disappears, whence a formula for $\sigma.s_{A_1}$. Comparing the latter with (9) and solving for N_d^1 , we obtain (1).

To obtain K_d^1 , consider the natural map

$$T_v \rightarrow f^*(T_{\mathbb{P}^2}).$$

Comparing with (6), we see that this map vanishes (simply) precisely at the singular points of fibres of π plus the preimages of cusps. Consequently

$$\begin{aligned} K_d^1 + 12\omega + T(1) &= c_2(f^*T_{\mathbb{P}^2} \otimes T_v^*) \\ &= 3N_d^1 + 3d\omega + 3T(d_1) - T(1), \end{aligned}$$

which yields (3). Finally (8) and (3) yield (5).

§2. Rationals.

We now consider analogous questions for rational curves. As above, and as in [R2], the idea is to study the fibred surface $\pi : X \rightarrow B$ corresponding to the 1-parameter family of rational curves of degree d through a set of general points $A_1, \dots, A_{3d-2} \in \mathbb{P}^2$. X may be realized in many ways as a blowup of a geometrically ruled surface. One such way, $b : X \rightarrow X' = \mathbb{P}(E)$ is the blowing down of the sum R of all fibre components not meeting $s_1 = s_{A_1}$. Now we have

$$s_1^2 = -m < 0,$$

where by [R2] we have

$$2m = \sum_{d_1+d_2=d} N_{d_1}^0 N_{d_2}^0 d_1 d_2 \binom{3d-4}{3d_1-2}$$

hence we may assume E has a nonvanishing section $\mathcal{O}$ with $\mathbb{P}(\mathcal{O}) = s_1 \subset X'$ and $M^* = E/\mathcal{O}$ of degree $-m$. In fact as X' possesses another section s_{A_2} disjoint from s_{A_1} , clearly $E = \mathcal{O} \oplus M^*$, but we won't need this here. Now let $\pi^* E^* \rightarrow \mathcal{O}_{X'}(1)$ be the canonical invertible quotient (in the antiGrothendieck sense). Then the composite $\pi^* M \rightarrow \mathcal{O}_{X'}(1)$ vanishes precisely on s_1 , hence

$$c_1(\mathcal{O}_{X'}(1)) \sim s_1 + m f', \quad f' = \text{fibre of } \pi',$$

hence

$$c_1(\mathcal{O}_X(1)) := c_1(b^* \mathcal{O}_{X'}(1)) \sim s_1 + m f \quad f = \text{fibre of } \pi.$$

Consequently, letting T_v as above be the vertical tangent sheaf of X , we have

$$c_1(T_v) = 2s_1 + m f - R.$$

As above, the natural map

$$T_v \rightarrow f^* T_{\mathbb{P}^2}$$

vanishes precisely on singular points of fibres and cusp preimages, hence

$$\begin{aligned} \sum N_{d_1}^0 N_{d_2}^0 d_1 d_2 \binom{3d-2}{3d_1-1} + K_d^0 &= c_2(f^* T_{\mathbb{P}^2} \otimes T_v^*) \\ &= 3N_d^0 - 3dm + 3f^* \mathcal{O}_{\mathbb{P}^2}(1).R + R^2 \end{aligned}$$

Simplifying, we obtain (2). For instance, $K_3^0 = 24$, as is well known [KS].

Finally, note that the ramification divisor of $f : X \rightarrow \mathbb{P}^2$, i.e. the degeneracy locus of the differential $T_X \rightarrow f^* T_{\mathbb{P}^2}$, consists now precisely of the cuspidal fibres plus the sections $s_{A_1}, \dots, s_{A_{3d-2}}$. Consequently

$$-2s_1 - m f + R + (2g - 2)f + 3f^* \mathcal{O}(1) \sim \sum s_{A_i} + K_d^0 f, g = g_d^0.$$

Hence taking dot product with s_1 we get

$$m + 2g - 2 = -m + K_d^0,$$

which yields (4).

References.

- [DH] S. Diaz, J. Harris: ‘Geometry of Severi varieties’. Trans. AMS. **309** (1988), 1–34.
- [F] W. Fulton: ‘Intersection theory’. Springer.
- [KS] S. Kleiman, R. Speiser: ‘Enumerative geometry of cuspidal plane cubics’. CMS Proc. **6** (1986), 227–268.
- [P] R. Pandharipande: ‘Getzler’s formula’, preprint.
- [R1] Z. Ran: ‘Enumerative geometry of singular plane curves’. Invent. Math. **97**, 447–465 (1989).
- [R2] _____: ‘Bend, break and count’, preprint (Duke eprint 9704004).